

Document details

< Back to results | 1 of 1

Export Download Print E-mail Save to PDF Add to List More... >

Full Text View at Publisher

Scientific Reports Open Access
Volume 8, Issue 1, 1 December 2018, Article number 4202

Polygonumins A, a newly isolated compound from the stem of Polygonum minus Huds with potential medicinal activities (Article)

Ahmad, R.^a, Sahidin, I.^b, Taher, M.^c, Low, C.^a, Noor, N.M.^a, Sillapachaiyaporn, C.^d, Chuchawankul, S.^e, Sarachana, T.^d, Tencomnao, T.^d, Iskandar, F.^f, Rajab, N.F.^f, Baharum, S.N.^{ad}✉

^aInstitute of Systems Biology, Universiti Kebangsaan Malaysia, Bangi, Selangor, Malaysia
^bLaboratory of Natural Products Chemistry, Faculty of Pharmacy, Universitas Halu Oleo, Kendari, Southeast Sulawesi, Indonesia
^cDepartment of Pharmaceutical Technology, Kulliyah of Pharmacy, International Islamic University of Malaysia, Jalan Istana, Kuantan, Pahang, Malaysia

View additional affiliations

Abstract

View references (51)

Polygonumins A, a new compound, was isolated from the stem of Polygonum minus. Based on NMR results, the compound's structure is identical to that of vanicoside A, comprising four phenylpropanoid ester units and a sucrose unit. The structure differences were located at C-3'''. The cytotoxic activity of polygonumins A was evaluated on several cancer cell lines by a cell viability assay using tetrazolium dye 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT). The compound showed the highest antiproliferative (p < 0.05) activities against K562 (Human Leukaemia Cell Line), MCF7 (Human breast adenocarcinoma cell line), and HCT116 (Colorectal cancer cells) cells. Cytotoxic studies against V79-4 cells were carried out and showed that polygonumins A was toxic at 50 µg/ml, suggesting that this compound may be used as an anticancer drug without affecting normal cells. Polygonumins A also showed promising activity as an HIV-1 protease inhibitor with 56% relative inhibition. Molecular docking results indicated that the compound possesses high binding affinity towards the HIV protease over the low binding free energy range of -10.5 to -11.3 kcal/mol. P. minus is used in Malaysian traditional medicine for the treatment of tumour cells. This is the first report on the use of P. minus as an HIV-1 protease inhibitor. © 2018 The Author(s).

ISSN: 20452322
Source Type: Journal
Original language: English
DOI: 10.1038/s41598-018-22485-5
Document Type: Article
Publisher: Nature Publishing Group

References (51)

View in search results format >

All Export Print E-mail Save to PDF Create bibliography

1 Poonthananiwatkul, B., Lim, R.H.M., Howard, R.L., Pibanpaknatee, P., Williamson, E.M.
Traditional medicine use by cancer patients in Thailand
(2015) *Journal of Ethnopharmacology*, 168, pp. 100-107. Cited 13 times.
www.elsevier.com/locate/jethpharm
doi: 10.1016/j.jep.2015.03.057
View at Publisher

Metrics ?

0 Citations in Scopus
0 Field-Weighted Citation Impact

PlumX Metrics
Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:
Set citation alert >
Set citation feed >

Related documents

GC-MS/Olfactometric characterisation and aroma extraction dilution analysis of aroma active compounds in Polygonum minus essential oil
Rusdi, N.A. , Goh, H.-H. , Baharum, S.N.
(2016) *Plant OMICS*
Volatile profiling of aromatic traditional medicinal plant, polygonum minus in different tissues and its biological activities
Ahmad, R. , Baharum, S.N. , Bunawan, H.
(2014) *Molecules*
A recent review on phytochemical constituents and medicinal properties of kesum (Polygonum minus Huds.)
Vikram, P. , Chiruvella, K.K. , Ripain, I.H.A.
(2014) *Asian Pacific Journal of Tropical Biomedicine*

View all related documents based

- 2 Li, X.-Z., Zhang, S.-N., Liu, S.-M., Lu, F.
Recent advances in herbal medicines treating Parkinson's disease
(2013) *Fitoterapia*, 84 (1), pp. 273-285. Cited 56 times.
doi: 10.1016/j.fitote.2012.12.009
[View at Publisher](#)
-
- 3 Hassan, W.W.
(2007) *Healing Herbs of Malaysia. Federal Land Development Authority (FELDA)*. Cited 2 times.
Kuala Lumpur
-
- 4 Hunter, M.
Australian kesom oil - A new essential oil for the flavour and fragrance industry
(1996) *Agro Food Industry Hi-Tech*, 7 (5), pp. 26-28. Cited 5 times.
-
- 5 Miyazawa, M., Tamura, N.
Components of the essential oil from sprouts of *Polygonum hydropiper* L. ('Benitade')
(2007) *Flavour and Fragrance Journal*, 22 (3), pp. 188-190. Cited 22 times.
doi: 10.1002/ffj.1779
[View at Publisher](#)
-
- 6 Vimala, S., Adenan, M.I., Ahmad, A.R., Shahdan, R.
(2003) *Nature's Choice to Wellness, Antioxidant Vegetables/ulam*. Cited 21 times.
Forest Research Institute Malaysia
-
- 7 Burkill H., 2 (Kuala Lumpur: Ministry of Agriculture & Cooperatives, 1966).
-
- 8 Hassim, N., Markom, M., Anuar, N., Dewi, K.H., Baharum, S.N., Mohd Noor, N.
Antioxidant and antibacterial assays on polygonum minus extracts: Different extraction methods ([Open Access](#))
(2015) *International Journal of Chemical Engineering*, 2015, art. no. 826709. Cited 2 times.
<http://www.hindawi.com/journals/ijce/>
doi: 10.1155/2015/826709
[View at Publisher](#)
-
- 9 George, A., Chinnappan, S., Chintamaneni, M., Kotak C, V., Choudhary, Y., Kueper, T., Radhakrishnan, A.K.
Anti-inflammatory effects of *Polygonum minus* (Huds) extract (Lineminus™) in in-vitro enzyme assays and carrageenan induced paw edema
(2014) *BMC Complementary and Alternative Medicine*, 14 (1), art. no. 355. Cited 10 times.
<http://www.biomedcentral.com/1472-6882/14/355>
-
- 10 Ali, A.M.
Antiviral and cytotoxic activities of some plants used in Malaysian indigenous medicine
(1996) *Pertanika Journal of Tropical Agricultural Science*, 19 (2-3), pp. 129-136. Cited 47 times.
-

Find more related documents in
Scopus based on:

Authors >

- 11 Qader, S.W., Abdulla, M.A., Chua, L.S., Sirat, H.M., Hamdan, S.
Pharmacological mechanisms underlying gastroprotective activities of the fractions obtained from *Polygonum minus* in sprague dawley rats

(2012) *International Journal of Molecular Sciences*, 13 (2), pp. 1481-1496. Cited 19 times.
<http://www.mdpi.com/1422-0067/13/2/1481/pdf>
doi: 10.3390/ijms13021481

[View at Publisher](#)

- 12 Azlim Almey, A.A., Ahmed Jalal Khan, C., Syed Zahir, I., Mustapha Suleiman, K., Aisyah, M.R., Kamarul Rahim, K.
Total phenolic content and primary antioxidant activity of methanolic and ethanolic extracts of aromatic plants' leaves

(2010) *International Food Research Journal*, 17 (4), pp. 1077-1084. Cited 64 times.
[http://www.ifrj.upm.edu.my/17%20\(04\)%202010/\(28\)%20IFRJ-2010-035%20Kamarul\[1\].pdf](http://www.ifrj.upm.edu.my/17%20(04)%202010/(28)%20IFRJ-2010-035%20Kamarul[1].pdf)

- 13 Huda-Faujan, N., Noriham, A., Norrakiah, A.S., Babji, A.S.
Antioxidant activity of plants methanolic extracts containing phenolic compounds

(2009) *African Journal of Biotechnology*, 8 (3), pp. 484-489. Cited 135 times.
<http://www.academicjournals.org/AJB/PDF/pdf2009/4Feb/Huda-Faujan%20et%20al.pdf>

[View at Publisher](#)

- 14 Maizura, M., Aminah, A., Aida, W.M.W.
Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma longa*) extract
(2011) *International Food Research Journal*, p. 18.

- 15 Baharum, S.N., Bunawan, H., Ghani, M.A., Wan Aida Wan Mustapha, Noor, N.M.
Analysis of the chemical composition of the essential oil of *Polygonum minus* Huds. Using two-dimensional gas chromatography-time-of-flight mass spectrometry (GC-TOF MS)

(2010) *Molecules*, 15 (10), pp. 7006-7015. Cited 46 times.
<http://www.mdpi.com/1420-3049/15/10/7006/pdf>
doi: 10.3390/molecules15107006

[View at Publisher](#)

- 16 Ahmad, R., Baharum, S.N., Bunawan, H., Lee, M., Noor, N.M., Rohani, E.R., Ilias, N., (...), Zin, N.M.
Volatile profiling of aromatic traditional medicinal plant, *Polygonum minus* in different tissues and its biological activities

(2014) *Molecules*, 19 (11), pp. 19220-19242. Cited 12 times.
<http://www.mdpi.com/1420-3049/19/11/19220/pdf>
doi: 10.3390/molecules191119220

[View at Publisher](#)

- 17 Ahmad, R., Bunawan, H., Normah, M.N., Baharum, S.N.
Chemical composition in different tissues of *Polygonum minus* by using GC X GC-TOF MS and direct discrimination by multivariate analysis of fourier transform infrared spectroscopy data

(2016) *International Journal of Pharmacognosy and Phytochemical Research*, 8 (12), pp. 1986-1992.
<http://impactfactor.org/PDF/IJPPR/8/IJPPR,Vol8,Issue12,Article14.pdf>

- ☐ 18 Zimmermann, M.L., Sneden, A.T.
Vanicosides a and b, protein kinase c inhibitors from *polygonum pensylvanicum*
(1994) *Journal of Natural Products*, 57 (2), pp. 236-242. Cited 34 times.
doi: 10.1021/np50104a007
[View at Publisher](#)
-
- ☐ 19 Cordell, G.A., Beecher, C.W.W., Pezzuto, J.M.
Can ethnopharmacology contribute to the development of new anticancer drugs?
(1991) *Journal of Ethnopharmacology*, 32 (1-3), pp. 117-133. Cited 102 times.
doi: 10.1016/0378-8741(91)90110-Y
[View at Publisher](#)
-
- ☐ 20 Boik, J. (Oregon Medical Press, Princeton, MN, USA, 2001).
-
- ☐ 21 Efferth, T., Miyachi, H., Bartsch, H.
Pharmacogenomics of a traditional Japanese herbal medicine (Kampo) for cancer therapy
(2007) *Cancer Genomics and Proteomics*, 4 (2), pp. 81-91. Cited 34 times.
<http://cgpp.iijournals.org>
-
- ☐ 22 Dzoyem, J.P., Nkuete, A.H.L., Kuete, V., Tala, M.F., Wabo, H.K., Guru, S.K., Rajput, V.S., (...), Tan, N.-H.
Cytotoxicity and antimicrobial activity of the methanol extract and compounds from *Polygonum limbatum*
(2012) *Planta Medica*, 78 (8), pp. 787-792. Cited 41 times.
doi: 10.1055/s-0031-1298431
[View at Publisher](#)
-
- ☐ 23 Lin, Y.-S., Huang, L.-D., Lin, C.-H., Huang, P.-H., Chen, Y.-J., Wong, F.-H., Lin, C.-C., (...), Fu, S.-L.
In Vitro and in Vivo anticancer activity of a synthetic glycolipid as toll-like receptor 4 (TLR4) activator
(2011) *Journal of Biological Chemistry*, 286 (51), pp. 43782-43792. Cited 15 times.
<http://www.jbc.org/content/286/51/43782.full.pdf+html>
doi: 10.1074/jbc.M111.285171
[View at Publisher](#)
-
- ☐ 24 Zhu, L., Cao, X., Chen, W., Zhang, G., Sun, D., Wang, P.G.
Syntheses and biological activities of daunorubicin analogs with uncommon sugars
(2005) *Bioorganic and Medicinal Chemistry*, 13 (23), pp. 6381-6387. Cited 38 times.
doi: 10.1016/j.bmc.2005.06.053
[View at Publisher](#)
-
- ☐ 25 Cingi, M.R., De Angelis, I., Fortunati, E., Reggiani, D., Bianchi, V., Tiozzo, R., Zucco, F.
Choice and standardization of test protocols in cytotoxicology: A multicentre approach
(1991) *Toxicology in Vitro*, 5 (2), pp. 119-125. Cited 49 times.
doi: 10.1016/0887-2333(91)90031-8
[View at Publisher](#)
-

- 26 Shrivastava, R.J., Rispat, G.W., Chevalier, G., Massingham, A., Can, R.
The in vivo maximum tolerated dose be predicted using in vitro techniques? A working hypothesis
(1991) *ATLA*, 19, pp. 393-402. Cited 25 times.

- 27 Borges de Melo, E., da Silveira Gomes, A., Carvalho, I.
 α - and β -Glucosidase inhibitors: chemical structure and biological activity

(2006) *Tetrahedron*, 62 (44), pp. 10277-10302. Cited 341 times.
doi: 10.1016/j.tet.2006.08.055

[View at Publisher](#)

- 28 Datta, B.K., Datta, S.K., Khan, T.H., Kundu, J.K., Rashid, M.A., Nahar, L., Sarker, S.D.
Anti-cholinergic, cytotoxic and anti-HIV-1 activities of sesquiterpenes and a flavonoid glycoside from the aerial parts of *Polygonum viscosum*

(2004) *Pharmaceutical Biology*, 42 (1), pp. 18-23. Cited 20 times.
doi: 10.1080/13880200490504943

[View at Publisher](#)

- 29 Lin, H.-W., Sun, M.-X., Wang, Y.-H., Yang, L.-M., Yang, Y.-R., Huang, N., Xuan, L.-J., (...), Xiao, K.
Anti-HIV activities of the compounds isolated from *polygonum cuspidatum* and *polygonum multiflorum*

(2010) *Planta Medica*, 76 (9), pp. 889-892. Cited 28 times.
doi: 10.1055/s-0029-1240796

[View at Publisher](#)

- 30 Zhong, Y., Yoshinaka, Y., Takeda, T., Shimizu, N., Yoshizaki, S., Inagaki, Y., Matsuda, S., (...), Yamamoto, N.
Highly potent anti-HIV-1 activity isolated from fermented *Polygonum tinctorium* Aiton

(2005) *Antiviral Research*, 66 (2-3), pp. 119-128. Cited 13 times.
doi: 10.1016/j.antiviral.2005.02.003

[View at Publisher](#)

- 31 Kim, H.J., Woo, E.-R., Shin, C.-G., Hwang, D.J., Park, H., Lee, Y.S.
HIV-1 integrase inhibitory phenylpropanoid glycosides from *Clerodendron trichotomum*

(2001) *Archives of pharmacal research*, 24 (4), pp. 286-291. Cited 48 times.
doi: 10.1007/BF02975093

[View at Publisher](#)

- 32 Qader, S.W., Abdulla, M.A., Chua, L.S., Najim, N., Zain, M.M., Hamdan, S.
Antioxidant, total phenolic content and cytotoxicity evaluation of selected Malaysian plants

(2011) *Molecules*, 16 (4), pp. 3433-3443. Cited 64 times.
<http://www.mdpi.com/1420-3049/16/4/3433/pdf>
doi: 10.3390/molecules16043433

[View at Publisher](#)

- 33 Fan, P., Terrier, L., Hay, A.-E., Marston, A., Hostettmann, K.
Antioxidant and enzyme inhibition activities and chemical profiles of *Polygonum sachalinensis* F.Schmidt ex Maxim (Polygonaceae)

(2010) *Fitoterapia*, 81 (2), pp. 124-131. Cited 56 times.
doi: 10.1016/j.fitote.2009.08.019

[View at Publisher](#)

- 34 Saint-Cricq De Gaulejac, N., Glories, Y., Vivas, N.
Free radical scavenging effect of anthocyanins in red wines

(1999) *Food Research International*, 32 (5), pp. 327-333. Cited 72 times.
doi: 10.1016/S0963-9969(99)00093-9

[View at Publisher](#)

- 35 Atta-ur-Rahman, Choudhary, M.I.
Bioactive natural products as a potential source of new pharmacophores. A theory of memory

(2001) *Pure and Applied Chemistry*, 73 (3), pp. 555-560. Cited 98 times.

[View at Publisher](#)

- 36 Papandreou, M.A., Dimakopoulou, A., Linardaki, Z.I., Cordopatis, P., Klimis-Zacas, D., Margarity, M., Lamari, F.N.
Effect of a polyphenol-rich wild blueberry extract on cognitive performance of mice, brain antioxidant markers and acetylcholinesterase activity

(2009) *Behavioural Brain Research*, 198 (2), pp. 352-358. Cited 134 times.
doi: 10.1016/j.bbr.2008.11.013

[View at Publisher](#)

- 37 Kapewangolo, P., Tawha, T., Nawinda, T., Knott, M., Hans, R.
Sceletium tortuosum demonstrates in vitro anti-HIV and free radical scavenging activity

(2016) *South African Journal of Botany*, 106, pp. 140-143. Cited 2 times.
<http://www.elsevier.com>
doi: 10.1016/j.sajb.2016.06.009

[View at Publisher](#)

- 38 Thayil, Seema, M., Thyagarajan, S.P.
Pa-9: A flavonoid extracted from plectranthus amboinicus inhibits HIV-1 protease

(2016) *International Journal of Pharmacognosy and Phytochemical Research*, 8 (6), pp. 1020-1024. Cited 2 times.
<http://impactfactor.org/PDF/IJPPR/8/IJPPR,Vol8,Issue6,Article19.pdf>

- 39 Wang, C.R., Zhou, R., Ng, T.B., Wong, J.H., Qiao, W.T., Liu, F.
First report on isolation of methyl gallate with antioxidant, anti-HIV-1 and HIV-1 enzyme inhibitory activities from a mushroom (Pholiota adiposa)

(2014) *Environmental Toxicology and Pharmacology*, 37 (2), pp. 626-637. Cited 14 times.
doi: 10.1016/j.etap.2014.01.023

[View at Publisher](#)

- 40 Razzaghi-Asl, N., Sepehri, S., Ebadi, A., Miri, R., Shahabipoura, S.
Effect of biomolecular conformation on docking simulation: A case study on a potent HIV-1 protease inhibitor

(2015) *Iranian Journal of Pharmaceutical Research*, 14 (3), pp. 785-802. Cited 2 times.
http://ijpr.sbmu.ac.ir/pdf_1674_c9745060f94d415c8778c1b925bf335c.html

- 41 Bunawan, H., Choong, C.Y., Md-Zain, B.M., Baharum, S.N., Noor, N.M.
Molecular systematics of polygonum minus huds. Based on ITS sequences

(2011) *International Journal of Molecular Sciences*, 12 (11), pp. 7626-7634. Cited 7 times.
<http://www.mdpi.com/1422-0067/12/11/7626/pdf>
doi: 10.3390/ijms12117626

[View at Publisher](#)

- ☐ 42 Ellman, G.L., Courtney, K.D., Andres Jr., V., Featherstone, R.M.
A new and rapid colorimetric determination of acetylcholinesterase activity

(1961) *Biochemical Pharmacology*, 7 (2), pp. 88-90,IN1-IN2,91-95. Cited 15368 times.

[View at Publisher](#)

-
- ☐ 43 Oyaizu, M.
Studies on products of browning reaction prepared from glucoseamine
(1986) *Japanese JNutr*, p. 44.

-
- ☐ 44 Wootton-Beard, P.C., Moran, A., Ryan, L.
Stability of the total antioxidant capacity and total polyphenol content of 23 commercially available vegetable juices before and after in vitro digestion measured by FRAP, DPPH, ABTS and Folin-Ciocalteu methods

(2011) *Food Research International*, 44 (1), pp. 217-224. Cited 176 times.
doi: 10.1016/j.foodres.2010.10.033

[View at Publisher](#)

-
- ☐ 45 Abe, K., Matsuki, N.
Measurement of cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction activity and lactate dehydrogenase release using MTT

(2000) *Neuroscience Research*, 38 (4), pp. 325-329. Cited 134 times.
doi: 10.1016/S0168-0102(00)00188-7

[View at Publisher](#)

-
- ☐ 46 Trott, O., Olson, A.J.
Software news and update AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading

(2010) *Journal of Computational Chemistry*, 31 (2), pp. 455-461. Cited 4680 times.
<http://www3.interscience.wiley.com/cgi-bin/fulltext/122439542/PDFSTART>
doi: 10.1002/jcc.21334

[View at Publisher](#)

-
- ☐ 47 Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., Ferrin, T.E.
UCSF Chimera - A visualization system for exploratory research and analysis

(2004) *Journal of Computational Chemistry*, 25 (13), pp. 1605-1612. Cited 12082 times.
doi: 10.1002/jcc.20084

[View at Publisher](#)

-
- ☐ 48 Takasaki, M., Konoshima, T., Kuroki, S., Tokuda, H., Nishino, H.
Cancer chemopreventive activity of phenylpropanoid esters of sucrose, vanicoside B and lapathoside A, from *Polygonum lapathifolium*

(2001) *Cancer Letters*, 173 (2), pp. 133-138. Cited 41 times.
doi: 10.1016/S0304-3835(01)00670-X

[View at Publisher](#)

-
- ☐ 49 Kiem, P.V., Nhiem, N.X., Cuong, N.X., Hoa, T.Q., Huong, H.T., Huong, L.M., Minh, C.V., (...), Kim, Y.H.
New phenylpropanoid esters of sucrose from *Polygonum hydropiper* and their antioxidant activity

(2008) *Archives of Pharmacal Research*, 31 (11), pp. 1477-1482. Cited 16 times.
doi: 10.1007/s12272-001-2133-y

[View at Publisher](#)

□ 50 Kawai, Y., Kumagai, H., Kurihara, H., Yamazaki, K., Sawano, R., Inoue, N.

β-Glucosidase inhibitory activities of phenylpropanoid glycosides, vanicoside A and B from *Polygonum sachalinense* rhizome

(2006) *Fitoterapia*, 77 (6), pp. 456-459. Cited 16 times.
doi: 10.1016/j.fitote.2006.05.008

[View at Publisher](#)

□ 51 Kumagai, H., Kawai, Y., Sawano, R., Kurihara, H., Yamazaki, K., Inoue, N.

Antimicrobial substances from rhizomes of the giant knotweed *Polygonum sachalinense* against the fish pathogen *Photobacterium damsela* subsp. *piscicida*

(2005) *Zeitschrift fur Naturforschung - Section C Journal of Biosciences*, 60 (1-2), pp. 39-44. Cited 12 times.

🔍 Baharum, S.N.; Institute of Systems Biology, Universiti Kebangsaan Malaysia, Bangi, Selangor, Malaysia;
email:nataqain@ukm.edu.my

© Copyright 2018 Elsevier B.V., All rights reserved.

[< Back to results](#) | 1 of 1

[^ Top of page](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語に切り替える](#)
[切换到简体中文](#)
[切换到繁體中文](#)
[Русский язык](#)

Customer Service

[Help](#)
[Contact us](#)

ELSEVIER

[Terms and conditions](#) [Privacy policy](#)

Copyright © 2018 Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

Cookies are set by this site. To decline them or learn more, visit our [Cookies page](#).

 RELX Group™